

GENERAL INDEX TO VOLUME XXXI

New scientific names of plants and the final members of new combinations are printed in bold-face type; synonyms and page numbers having reference to figures and plates, in *italics*; and previously published and all other matter, in ordinary type.¹

A

- Acerates Florida*, 368
 Adaptation in yeasts, 219
Ageratum lineare, 265
 Alkali Ridge, Utah, prehistoric maize collection from, 345, 350, 352
 Alleles, mating type, Instability of the, in *Saccharomyces*, 203
Amauroascus, 186
 Ammerman, Elizabeth, see Elizabeth Ammerman Baltzer
 Amphiploidy, see Polyploidy
 Anderson, Edgar. Cytological observations on *Tripsacum dactyloides*, 317; Homologies of the ear and tassel in *Zea Mays*, 325; Maiz reventador, 301; Notes on variation in *Tithonia tubaeformis*, 239; The sources of effective germ-plasm in hybrid maize, 355; Two collections of prehistoric corn tassels from southern Utah, 345; and Dorothy Schregardus. A method for recording and analyzing variations of internode pattern, 241
 Araceae of Panama, 1
 Arachniotus, the genus, 173; cytological study of, 174; history of, 173; systematic position of, 188
Arachniotus aureus, 173, 174, 186; *canadensis*, 173, 174, 186; *citrinus*, 174, 186; *ruber*, 173, 174, 186; *trachyspermus*, 174, 186; *trispurus*, 173, 174, 187, 190, 192, 194, 196, 198, development and systematic position of, The, 173
 Archeological collections of corn tassels, 304, 335, 345, 350, 352
 Asclepiadaceae from tropical America, Miscellaneous new, 235
 Asclepiads, Notes on some North American, 363
Asclepias, 363; *decumbens*, 368; *fascicularis*, 364, 365; *Florida*, 368; *galioides*, 365; *incarnata*, 363; *Linaria*, 364; *mexicana*, 363; *revoluta*, 368; *Rolfii*, 368; *subverticillata*, 365; *tuberosa*, 366; ssp. *interior*, 368, 370, ssp. *tuberosa*, 368, 370, ssp. *Rolfii*, 368; *tuberosa*, 366, 368, var. *decumbens*, 367, var. *flexuosa*, 368; *verticillata*, 365, var. *subverticillata*, 365

- Aspergillaceae*, 188
Axillaria, 382

B

- Baltzer, Elizabeth Ammerman. A monographic study of the genus *Palafoxia* and its immediate allies, 249
Bromeliaceae of Panama, 73
 Butterflyweed, 366

C

- Camphor, effect of, on copulation in *Saccharomyces cerevisiae*, 203, 205, 206
Campydomum, 382
 Chiasma frequencies in *Tripsacum*, 320
 Chromosome numbers in: *Coelorachis*, 321; *Tripsacum dactyloides*, 319; *Manisuris*, 321; *Polygonatum*, 377, 403; *Rottboellia*, 321; *Zea*, 319
 Clines, 369
Commelinaceae of Panama, 138
Convallaria, 382; *angustifolia*, 393; *biflora*, 393; *canaliculata*, 393; *commutata*, 401; *gigantea*, 402; *hirta*, 412; *parviflora*, 393; *Polygonatum*, 382; *pubescens*, 383
 Copulation: in *Arachniotus trispurus*, 180, 182; in *Saccharomyces cerevisiae*, 203, effect of camphor on, 203, 205, 206
 Corn tassels, prehistoric, from southern Utah, Two collections of, 345
 Corn, see Maize
 Crozier formation in *Arachniotus*, 173, 188
Ctenomyces, 186; *serratus*, 188
Cynanchum cubense, 236; *Haughtii*, 235; *Marsdenioides*, 235; *peraffine*, 236
 Cytological observations: on *Arachniotus trispurus*, 174; on *Polygonatum*, 377; on *Tripsacum dactyloides*, 317

D

- Dermatophytes*, 188
 Diakinesis, *Tripsacum dactyloides* mother cells at, 320
 Diploid yeast cultures: adaptation in, as contrasted with haploid, 231; criteria for distinguishing, 209

E

- Ear and tassel in *Zea Mays*, Homologies of the, 325

¹ For Woodson & Schery's "Flora of Panama" only the plant families and new entities are included in the *ANNALS* Index, since a complete Index will be appended at the end of each volume of the "Flora."

Enzymatic adaptation in yeasts, 219
 Eriocaulaceae of Panama, 65
 Euasclepias, 363
 Euchlaena, 325; mexicana, 307
 Evallaria, 382

F

Favotrichum ochraceum, 188
 Fernald's view of Asclepias tuberosa problem, 366
 Fertilization in Arachniotus trisporus, 181
 Flora of Panama, Part II, Fasc. 3, 1
 Florestina, 254; callosa, 255
 Frequency distributions for two prehistoric collections of corn, 348, 349

G

Gaillardia lanceolata, 263
 Galactose fermentation in yeasts, 219
 Germ-plasm in hybrid maize, The sources of effective, 355
 Gymnosaceae, 173, 188; crozier formation in, 173
 Gymnosiscus, 186; aureus, 173, 186; candidus, 173, 186; Reessii, 188; ruber, 173, 186; sudans, 173, 174, 177, 187

H

Haploid yeast cultures, criteria for distinguishing, 209
 Heiser, Charles B., Jr. Monograph of Psilostrophe, 279
 Hershey's story of Lancaster Surecropper corn, 358
 Homologies of the ear and tassel in Zea Mays, 325
 Hotson's strain of Arachniotus trisporus, 173, 174, 188, 200
 Hybrid maize, The sources of effective germ-plasm in, 355
 Hymenopappus integrifolius, 263

I

Inflorescence in Zea Mays; main features of male, 327, 346, 346; female, 325
 Instability of the mating type alleles in Saccharomyces, 302, under selection pressure, 207
 Internode pattern: A method for recording and analyzing variations of, 241; of Salvia, 243, 245; of Tradescantia, 243, 244; of Tripsacum dactyloides, 245, 246

J

Jalisco: maize grown in, 301; ears of maíz reventador from, 312, 314; Tithonia

tubaeformis in, 239
 Kelly, Dr. Isabel, 301
 Kinetics of enzymatic adaptation in genetically homogeneous and heterogeneous populations of yeast, A comparison of the, 219

L

Lancaster Surecropper corn, history of, 358
 Lemnaceae of Panama, 60
 Liliaceous genus Polygonatum in North America, The, 373
 Lindegren, Gertrude, Carl C. Lindegren and: Instability of the mating type alleles in Saccharomyces, 203
 Lindegren, Carl C.: and Gertrude Lindegren. Instability of the mating type alleles in Saccharomyces, 203; S. Spiegelman and. A comparison of the kinetics of enzymatic adaptation in genetically homogeneous and heterogeneous populations of yeasts, 219
 Lomaxeta, 262; verrucosa, 263

M

Maíz reventador, 301, 305, 312, 314; distribution of, 306, 309; uses of, 311; varieties mixed with, 314
 Maiz, see Maize
 Maize: Basket Maker, 349; Bearpaw Dent, 335, 342; breeding, 333, 337, 355; ear-tassel correlation in, 326, 329, 342; genetic background of, 330; germ-plasm, Sources of effective, in hybrid, 355; history of, in the Southwest, 349; Homologies of ear and tassel in, 325; hybrid, 307, 355; inbred lines of, 326; Inbred "Os 420", 335, 342; Krug, 356; Lancaster Surecropper, 342, 355; maíz chapolote, 302, 312; Maíz reventador, 301; maíz rosquera, 301; open-pollinated, 355, Yellow Dent, 335, 342, 355; Mexican, 301, 329, 350, 352; Pueblo, 342; plant color, 303; prehistoric corn tassels from southern Utah, Two collections of, 345, 350, 352; Reid, 356; tassel, morphology of, 327, 327, score sheet showing technical details of, 347; Yellow Dent, 335, 342, 357
 Manisuris, chromosome numbers of, 321
 Matelea serpens, 236
 Mayacaceae of Panama, 62
 Media: for growing Arachniotus trisporus, 184; for growing Saccharomyces, 222
 Megatrachophyton equinum, 188; ferrugineum, 188

- Method for recording and analyzing variations of internode pattern, A, 241
- Mexican corn, 301, 329; introduction in American Southwest, 350
- Mexico: maize in, 301; distribution of maíz reventador in, 309, map, showing, 306
- Microsporon, 188; Audouini, 188; lanosum, 188
- Milkweeds, 363
- Moldenke, Harold N. Eriocaulaceae of Panama, 65
- Monograph of Psilostrophe, 279
- Monographic study of the genus Palafoxia and its immediate allies, A, 249
- Myxotrichum uncinatum, 188
- N
- Nannizzi's strain of Arachniotus trisporus, 174, 187, 188, 200
- Natural selection, mathematical theory of, 223; applied to yeasts, 226
- Notes on some North American Asclepiads, 363
- Notes on variation in Tithonia tubaeformis, 239
- O
- Ownbey, Ruth Peck. The Liliaceous genus Polygonatum in North America, 373
- Othake, 249, 253; callosum, 255, distribution of, 270; canescens, 257, distribution of, 272; Hookerianum, 261, distribution of, 274; Hookerianum, 260; Lindenii, 258, distribution of, 274; macrolepis, 258; maximum, 261; Reverchonii, 259, distribution of, 274; robustum, 257; roseum, 256, distribution of, 272, var. robustum, 257, distribution of, 272; sphacelatum, 259, distribution of, 274; tenuifolium, 255, 262; texanum, 257, distribution of, 274, var. macrolepis, 258, distribution of, 274
- P
- Palafoxia, A monographic study of Palafoxia and its immediate allies, 249
- Palafoxia, 249, 264; arenaria, 266; callosa, 255; fastigiata, 263; Feayi, 267, distribution of, 276; Hookeriana, 260, 261, var. subradiata, 260; integrifolia, 263; leucophylla, 266; Lindenii, 258; linearis, 265, 270, distribution of, 276, var. arenicola, 266, var. gigantea, 266, distribution of, 276, var. leucophylla, 266, distribution of, 276; linearis, 266; Texana, 257, 261
- Palafoxia, 253, 262
- Paleolaria, 262, 264; carnea, 265; fastigiata, 263
- Panama, Flora of, Part II, Fasc. 3, 1
- Pathogenicity of Arachniotus trisporus, 185
- Peabody Museum of Harvard University, prehistoric corn tassels in, 345
- Periballanthus, 382
- Pinole, maíz reventador used in, 311
- Pioneer Hi-Bred Corn Company, tests with inbred lines of maize by, 333
- Pipitillo, 301
- Polygonatum, The Liliaceous genus, in North America, 373
- Polygonatum, 382; angustifolium, 393; biflorum, 393, distribution of, 395, var. hebetifolium, 399, var. melleum, 399, var. necopinum, 401; biflorum, 383, 400, var. commutatum, 401, var. giganteum, 402, var. hebetifolium, 393, γ. latifolium, 383, var. ovatum, 402, var. ovatum subvar. ellipsoidale, 402, f. ramosum, 402, var. virginicum, 402; boreale, 384, var. australe, 384, var. multiflorum, 384; canaliculatum, 393, var. americanum, 402, var. americanum subvar. ellipticum, 402, β. giganteum, 402, var. giganteum, 402, var. oblongifolium, 402, var. ovatum, 402, f. ramosum, 402; cobrense, 392; commutatum, 401, distribution of, 404; commutatum var. lineamentosum, 402, var. ovatum, 402, f. ramosum, 402, var. virginicum, 402; cuneatum, 383; ellipticum, 402; Farwellii, 384; giganteum, 402; hebetifolium, 393, 400; birtum, 412; latifolium, 402, var. commutatum, 401; melleum, 399; multiflorum, β. americanum, 383, 393, 402, γ. biflora, 393, δ. canaliculata, 393, γ. giganteum, 402, ε. latifolia, 402, δ. latifolium, 383, β. pubescens, 383; officinale, 382; ovatum, 402; parviflorum, 393; pubescens, 383, distribution of, 385; pubescens var. australe, 384, var. boreale, 384, var. boreale subvar. australe, 384, var. cuneatum, 383, f. cuneatum, 384, f. fultius, 384, var. multiflorum, 384; virginicum, 402
- Polyploidy in Tripsacum dactyloides, 317; in Polygonatum, 377
- Polypteris, 249, 262; brasiliensis, 263; callosa, 255; Hookeriana, 260, 261; integrifolia, 262, 270, distribution of, 276; Lindenii, 258; macrolepis, 258; maxima, 261; Reverchonii, 259; robustum, 257; rosea, 256; Texana, 257
- Polypteris 254

- Pontederiaceae of Panama, 151
 Popcorn from maíz reventador, 301
 Populations of yeasts, 226
 Prehistoric corn tassels from southern Utah,
 Two collections of, 345, 350, 352
 Psilostrophe, Monograph of, 279
 Psilostrophe, 285; *Bakeri*, 288, 300, distribution of, 283; *biennis*, 293; *cerifera*, 293, var. *biennis*, 293; *Cooperi*, 287, 300, distribution of, 284; *divaricata*, 289, 290; *gnaphalodes*, 294, 300, distribution of, 283; *grandiflora*, 289, 292; *Hartmanii*, 290; *lanata*, 292; *pumila*, 288; *sparsiflora*, 289, 300, distribution of, 283; *sparsiflora*, 293; *tagetina*, 290, 300, distribution of, 284, var. *grandiflora*, 292, 300, distribution of, 284, var. *lanata*, 292, distribution of, 284, var. *sparsiflora*, 289; *Tagetinae*, 290, villosa, 293, 300, distribution of, 283
 Pueblo corn, 349
- Q
- Quantitative inheritance in maize, 326, 339
- R
- Rapateaceae of Panama, 71
 Reproduction, sexual, in *Arachniotus trisporus*, 180; ascus formation, 183, 198; copulation branches, 180, 182, 196; crozier formation, 183, 198; fertilization, 181
Riddellia, 285; *arachnoidea*, 294; *Cooperi*, 287; *gnaphalioides*, 294; *tagetina*, 290, var. *pumila*, 288, var. *sparsiflora*, 289
 Rosenbaum, Elisabeth Heuser. The development and systematic position of *Arachniotus trisporus*, 173
- S
- Saccharomyces cerevisiae*: Instability of the mating type alleles in, 203; galactozymase activity and natural selection in, 231
Sahuaripa, Relacion of (San Miguel de), manuscript copy of, 310
Salomonis, 382; *biflora*, 393; *cobrensis*, 392; *commutata*, 402
Salvia, internode pattern for species of, 245, 245
 Schery, Robert W., Robert E. Woodson, Jr. and. Flora of Panama, Part II, Fasc. 3, 1
 Schregardus, Dorothy, Edgar Anderson and. A method for recording and analyzing variations of internode pattern, 241
Sigillum, 382
Sipbyalis, 382
 Smith, Lyman B. Bromeliaceae of Panama, 73
 Sources of effective germ-plasm in hybrid maize, The, 355
 Spiegelman, S., and Carl C. Lindegren. A comparison of the kinetics of enzymatic adaptation in genetically homogeneous and heterogeneous populations of yeast, 219
 Standley, Paul C. Araceae of Panama, 1
Stevia, 253; *callosa*, 255; *linearis*, 265; *lavendulaefolia*, 265; *spbacelata*, 259
 Subspecies, three, of *Asclepias tuberosa*, 366
- T
- Tassels, maize: correlation between node condensation and row number, 333, 333; diagram showing main features of, 327, 340; morphology of, 326, 327; of maíz reventador, 308; prehistoric, from southern Utah, Two collections of, 345; profiles, correlation between ear shape and, 326, 329, 336, of four inbred strains, 332; score sheet of, 347; spikelet arrangement in, 328
 Temperature, optimum, for growth of *Arachniotus trisporus*, 185
 Teosinte, 307
Tithonia tubaeformis, Notes on variation in, 239
 Torula: transformation of *Saccharomyces* into, 203; *utilis*, 203
 Torulopsis, 203
Tradescantia, internode pattern for species of, 243, 244
Trichophyton, 188; *asteroides*, 188; *denticulatum*, 188; *felineum*, 188; *radiolatum*, 188
Trichophytoneae, 188
Tripsacum dactyloides: chiasma frequencies in, 320; chromosome numbers of, 319; Cytological observations on, 317; distribution of, 318; var. *occidentale*, 317; internode pattern of, 245, 246; pollen mother cells of, at diakinesis, 320
- U
- Utah, southern, Two collections of prehistoric corn tassels from, 345
- V
- Valonis' strain of *Arachniotus trisporus*, 173, 187, 200

Variation in *Tithonia tubaeformis*, Notes on, 239

Variations of internode pattern, A method for recording and analyzing, 241

W

Woodson, Robert E., Jr.: Miscellaneous new Asclepiadaceae from tropical America, 235; Notes on some North American Asclepiads, 363; Robert W. Schery, and collaborators. Flora of Panama, Part II, Fasc. 3, 1

X

Xyridaceae of Panama, 63

Y

Yeast, A comparison of the kinetics of enzymatic adaptation in genetically homogeneous and heterogeneous populations of, 219

Yeasts: effect of camphor on copulation of, 203, 205, 206, 212, 214, 216; haploid and diploid, criteria for distinguishing, 209, nuclear fusion in, 210; instability of mating type alleles in, 203

Z

Zea, chromosomes of, 319; Mays, Homologies of the ear and tassel in, 325

Zea Mays, see Maize

Zygosaccharomyces, 203